On the first day, January 29, thunder was heard from southward at intervals during the morning, and at 9^h 30^m a solar halo was seen. The sky became quite covered with heavy nimbus clouds soon after 13h, and at 14h 35m rain commenced to fall, lasting till 15^h 5^m, accompanied by occasional loud peals of thunder. The storm traveled from south to northnortheast passing slightly to the east of the observatory. The temperature began to fall before the sky was completely covered, and continued to do so until 14^h 30^m; at 13^h 50^m the wind veered from north to south, and was from southeast by south during the rain which fell from 14^h 30^m to 15^h 5^m.

On the second day the sky became covered with heavy nimbus clouds at 12^h0^m, and very heavy rain fell from 12^h 50^m to 13^h 10^m. On this occasion the storm traveled from southsouthwest to north-northeast, passing very nearly over the observatory; at the same time a second storm burst over the Ponce, a mountain 7 miles south by west of the observatory.

The fall of temperature during this storm was the greatest on record, viz, from 93.5° at 11^h 25^m to 72.0° at 12^h 58^m. Here also the fall commenced some minutes before the sky became covered.

The sunshine strip shows a continuous burned line up to 11^h 53^m, and a small spot at 12^h 0^m.

Between 11^h 25^m and 11^h 35^m, the wind veered from northnortheast to south, and was from the latter direction during the heavy rain which followed.

The registers are brought to the notice of meteorologists, with a view to obtaining further information on the effect of thunderstorms, and any such information will be much appreciated.

RECENT PAPERS BEARING ON METEOROLOGY.

W. F. R. PHILLIPS, in charge of Library, etc.

The subjoined list of titles has been selected from the contents of the periodicals and serials recently received in the library of the Weather Bureau. The titles selected are of papers or other communications bearing on meteorology or cognate branches of science. This is not a complete index of the meteorological contents of all the journals from which it has been compiled; it shows only the articles that appear to the compiler likely to be of particular interest in connection with the work of the Weather Bureau:

Journal de Physique. Paris. 3me série. Tome 9.

Brillouin, Marcel. Origine, variation et perturbations de l'électricité atmosphérique. P. 91.

Teisserenc de Bort. Étude de l'atmosphère dans la verticale

par cerfs-volants et ballons-sondes. P. 129. Scientific American. New York. Vol. 82.

Crafts, H. A. tains. P. 133.

Annalen der Physik.
Schwalbe, G. Snowfall and Water Supply of the Rocky Moun-

Leipzig. Vierte Folge. Band 1.

chwalbe, G. Uber die experimentalle Grundlage der Exner'schen theorie der Luftelectricität. P. 294.

Zeitschrift für Luftschiffahrt u. Physik der Atmosphäre. Berlin. 19 Jahrg.
Jacob, E. Forsetzung der Betrachtungen über eine kinetische
Theorie der Luftbewegungen. P. 5.
Nimfuhr, R. Flugtechnische Betrachtungen (Schluss). P. 14.

Nature. London. Vol. 61.

Dexter, E. G. Drunkenness and the Weather. P. 365.

Marconi, G. Wireless Telegraphy. P. 377.

Wood, R. W. Effects of Lightning on Electric Lamps. P. 391. ity and the Earth's Motion. P. 445.

W. E. Applied Meteorology. P. 448. ure. Paris. 28me Année.

La Nature. Paris. 28me Année.

Jaubert, J. L'Orage du 13 Février 1900. P. 210.

Naturwissenschaftliche Rundschau. Braunschweig. 15 Jahrg.

Fassig, O. L. Typen des März-Wetters in den Vereinigten Staaten. (Abstract American Journal Science). P. 94.

Das Wetter.

Wetter. Berlin. 17 Jahrg. Kasner C. Der Mistral. P. 25.

Meinardus, W. Ueber der Methoden der Maritimen Klimato-

Bornstein, R. Eine Verbesserung des telegraphischen Witterungsdienstes. P. 36.

Ciel et Terre. Bruxelles. 20me Année. Vander Linden, E. Prévision du temps pour une période de

plusieurs jours. P. 589.

Proceedings of the Royal Society. London. Vol. 66.

Sworn, S. A. Researches in Absolute Mercurial Thermometry. P. 86.

Gaea. Leipzig. 36 Jahrg.
Trabert, W. Die Bildung des Hagels. (Schluss). P. 207.
Symons's Meteorological Magaztne. London. Vol. 35.

— The Snow and Floods of February, 1900 [England]. P. 18.

Meteorologische Zeitschrift. Wien. Band 17

Less, E. Ueber den täglichen Gang der Sommerregen bei verschiedenen Wetterlagen. P. 49. Bergholz, P. Ueber Bildungsstätten, Bahnen und Zonen der

Orkane des Fernen Ostens. P. 71.

Woeikof, A. Arktis und Antarktis.

Woeikof, A. Al. v. Tillo. P. 79.

Hann, J. Klima von Ponta Delgada. P. 80.
 Hann, J. Ueber eine als möglich gedachte Ursache der Wirksamkeit des Hagelschiessens. P. 83.

Hann, J. Klimatabelle für Auckland (Neuseeland, Nordinsel). P. 84.

-Einfluss der Grossen Seen auf den Niederschlag. P. 87.

Kohlensäuregehalt der Luft auf dem Montblanc. P. 87

Neue tägliche Wetterkarten. P. 88. - Das Vorkommen von Jod in der Atmosphäre. P. 88.

MEXICAN CLIMATOLOGICAL DATA.

Through the kind cooperation of Senor Manuel E. Pastrana, Director of the Central Meteorologico-Magnetic Observatory, the monthly summaries of Mexican data are now communicated in manuscript, in advance of their publication in the Boletin Mensual. An abstract, translated into English measures, is here given, in continuation of the similar tables published in the Monthly Weather Review since 1896. The barometric means have not been reduced to standard gravity, but this correction will be given at some future date when the pressures are published on our Chart IV.

Mexican data for February, 1900

	Je.	ba- ter.	Ten	perat	ure.	tive dity.	ita-	Prevailing direction.		
Stations.	Altitude.	Mean	Max.	Min.	Меап.	Rela i	Precipit	Wind.	Cloud.	
	Feet.	Inch.	∘ <i>F</i> .	\circ_{F} .	∘F.	5	Inch.			
Colima	1,656		86.4	59.2						
Culiacán Rosales (E.										
d. S.)	112	29.76	86.0	56 1	71.4	64	3.22	w.		
Durango (Seminario).		24,00	77.0	28.4	54.5	47	0.94	sw.	sw.	
Leon (Guanajuato)	5, 934	24.28	79.0	36.1	59.4	44	0.04	sw.	s₩.	
Mazatlan		29.96	95.4	52.0	75.7	65	0.79	se.	е.	
Mexico (Obs. Cent.)		23.03	79.0	35.6	59.0	49	Т.	8.	w.	
Morelia (Seminario)		23.96	76.8	39.9	59.7	54		s₩.	BW.	
Puebla (Col. Cat.)		23.86	75.4	39.2	61.5	60	T.	ene., e.	8.	
Saltillo(Col. S. Juan).	5, 399	24.74	76.3	25.2	52.3	58	0.98	s.	sw.	
San Isidro (Hac. de							_			
Guanajuato)			69.4	56.8		• • • • • •	Т.	w.		
Silao		24.26	74.5	44.2	61.5	47	0,23	w .	w.	
Zapotlan	5,078	25.09	82.0	44.6	62.8	50	0.08	sw.	w.	

EARTHQUAKES AT CARSON CITY, NEV.

By Prof. Charles W. FRIEND.

According to a letter recently received from Prof. Charles W. Friend, the geographical position of his private observatory at Carson City, Nev., is latitude N. 39° 9′ 47.2″, and longitude W. 119° 45′ 42.9″. The altitude above sea level is 4,660 feet. The observatory is well furnished with a 5-inch refractor by Clark, a 3-inch transit by Troughton and Simms, a chronograph by Fauth, sideral clock, chronometers, sextants, and astro-photographic apparatus.

The seismograph is a duplex pendulum pattern from designs by Prof. J. A. Ewing. Meteorological records have been kept tridaily at 7 a. m., 2 p. m., and 9 p. m., Pacific standard time, since 1880. In the following list of earthquakes the scale of terms used in the 6th column is that known as the

Rossi-Forel scale, which reads as follows:

THE ROSSI-FOREL SCALE

I.—Microseismic shock, recorded by a single seismograph, or by seismographs of the same model, but not putting seismographs of different patterns in motion. Reported by experienced observers only. II.—Shocks recorded by several seismographs of different patterns.

Reported by a small number of persons who are at rest.

III.—Shocks reported by persons at rest. Duration or direction noted. IV.—Shocks reported by persons in motion. Shaking of movable objects, doors and windows, cracking of ceilings.

V.—Shock generally felt by every one; furniture shaken, some

VI.—General awakening of sleepers; general ringing of bells; swinging of chandeliers; stopping of clocks; visible swaying of trees; some persons run out of buildings.

VII.—Overturning of loose objects; fall of plaster; striking of church

bells; general fright; without damage to buildings.

VIII.—Fall of chimneys; cracks in walls of buildings. IX.-Partial or total destruction of some buildings.

X.—Great disasters; overturning of rocks; fissures in the surface of the earth; mountain slides.

Earth	quakes	observed	at	Carson	City,	Nev.
e, Pa- stand- rd.	on.	nsity, Forel			R	emark

		Haring	zuakes oo	servea	at Carson City, Nev.	
Year.	Month and date.	Time, Pa- cific stand- ard.	Motion.	Intensity, Rossi-Forel scale.	Remarks.	
1875	Jan. 24 Dec. 3	4.00 a.m. 3.00 p.m.	ne. sw.		One light and one sharp shock.	
1877 1881	July 9 Oct. 21 Nov. 9	11.00 p.m. 6.41 p.m. 10.08 a.m.	n. s. n. s. n. s.	:	Light. Do. Two light shocks. Sharp shock, lasting 2 seconds.	
1883	Inlv 1	3.00 a.m.			Light.	
1884 1887	Aug. 19 April 11 June 3	2.55 a. m. 2.10 p.m. 2.48 a. m.	nw. se. sw. ne.	VIII	Three light shocks. Principally vertical. Very severe, lasting 6 to 70 seconds, rotary motion preceded by a noise like thunder; stone and brick walls cracked, plaster shaken down, etc.	
1888	June 18 Jan. 29	1.20 a.m. 10.45 p.m.	sw. ne.		Two light shocks.	١,
	April 18 April 28 May 27	7.83 p.m. 8.47 p.m. 1.50 a.m.	sw. ne. s. n. s. n.	IV	Light. Lasting 5 to 6 seconds. Light, followed by quite heavy shock 20 seconds later.	1
1889	June 19 Oct. 15 Dec. 14	10.00 p.m. 4.80 a.m. 5.30 a.m.	s.n. e.w. e.w.	II II	Light. Do. Do.	1
1890 1892	April 24 Feb. 23	D. N. ¹ D. N. ¹		Į Į	Light (from seismometer.)	1
	Mar. 26	Bet. 7a.m.	} e. w.		Light tremors all day.	3
	April 19	2.51 a.m.	e. ₩.	VI	Gentle, but large movement, stopped sideral and meridian time clocks in observatory.	7 04 04 0
	April 21 April 21	9.44 a.m. 7.17 p.m.	e. w. e. w.	VI IV	observatory.	1
	April 23	5.30 p.m.	sw. ne.	II		1
,	April 29 May 28	4.08 p.m. D. N.	se. nw.	III	Light (from seismometer.)	1
	May 28 July 6 July 22	7.00 a.m. 6.50 a.m.	e. w. se. nw.	II	Light. Do.	!
1893	Mar. 2	12.05 a.m.	e. w.	11	Do.	1
	Mar. 2 Mar. 30	6.40 a.m. D. N. ¹	e.w. ne sw.	II	Do. Tremor (from seismometer.)	۱,
1894	Dec. 11 Nov. 10	8.10 p.m. 6.55 p m.	e. w. e. w.	II	Do. Light.	
109-8	Nov. 15	11.07 p.m.	e w.	I	Ďo,	-
	Nov. 15 Nov. 15	11.25 p. m. 12 midn't.	e. w.	II II	Do. Do.	1
	Nov. 18 Nov. 18	2.38 a.m. 2.40 a.m.	e. w. e. w.	I	Tremor (from selsmometer.) Do.	2
	Nov. 18	2.49 a.m.	e. w.	III	Sharp shock.	,
	Nov. 18 Nov. 18	5.15 a.m. 5.33 a.m.	e. w. e. w.	I	Tremor. Do.	ĺ
	Nov. 18 Nov. 21	7.22 a.m. D. N. ¹	e w.	I	Do. Tremor (from seismometer.)	
	NOV. 24	10.03 p.m.	sw. ne.	II	Light.	
	Nov. 24 Dec. 4	11.22 p.m. 9.89 p.m.	sw. ne.	III I	Sharp shock. Tremor.	
1896	Dec. 18 Jan. 25	9.09 p.m. 4.45 a.m.	sw. ne. e. w.	111	Sharp shock. Light.	l
1090	Jan. 25	4.46 a.m.	e. w.		Do.	
	Jan. 25 Jan. 27	5.02 a.m. 7.59 a.m.	e. w. s. n.		Do. Light, and a number of light tremors.	1
	Jan. 27 Jan. 27	8.34 a.m. 11.04 a.m.	w.e. sw.ne	III	Do. Do.	4
	Jan. 27	11.19 a.m.	sw. ne.	I	Do.	۱,
	Jan. 27 Jan. 27	1.01 p.m. 6.82 p.m.	sw. ne.	IV II	Do. Do.	
	Mar. 19 Mar. 20	4.01 a.m. 11.25 p.m.	• • • • • • • • • • • • • • • • • • • •		Light, Do.	1
1897	May 15	11.05 p.m.	sw. ne.	ΪΪΪ	. 20.	1
	May 21 June 20	1.50 p.m. 12.15 p.m.	sw. ne.	III	Stopped four clocks in United States Government building.	
	July 5 July 11	6.52 p.m. 12.15 a.m.			Tremor. Do.	1
4000	Oct. 14	10.30 a.m.	sw. ne.	ΪΪ	, 20.	į.
1898	Mar. 13 Mar. 30	7.34 a.m. 11.45 p.m.	sw. ne. sw. ne	11 1V		1
	1	i -		l		ı,

1 During night.

OBSERVATIONS AT HONOLULU.

Through the kind cooperation of Mr. Curtis J. Lyons, Meteorologist to the Government Survey, the monthly report of meteorological conditions at Honolulu is now made partly in accordance with the new form, No. 1040, and the arrangement of the columns, therefore, differs from those previously published.

Meteorological observations at Honolulu, February, 1900.

The station is at 21° 18' N., 15° 50' W.

Pressure is corrected for temperature and reduced to sea level, and the gravity correction, —0.06, has been applied.

The average direction and force of the wind and the average cloudiness for the whole day are given unless they have varied more than usual, in which case the extremes are given. The scale of wind force is 0 to 12, or Beaufort scale. Two directions of wind, or values of wind force or amounts of cloudiness, connected by a dash, indicate change from one to the other.

The rainfall for twenty-four hours has always been measured at 10:29 p. m., not 1 p. m., Greenwich time, on the respective dates.

The rain gaze, 8 inches in diameter. Is 1 foot above ground. Thermometer, 9 feet above ground. Ground is 43 feet, and the barometer 50 feet above sea level.

Date.	evel.	Tempera- ture.		During twenty-four hours preceding 1 p.m Green- wich time, or 2:29 a.m., Honolulu time.									98.
	ses le			Tempera- ture.		Means.		Wind.		-ipnc	Sea-level pressures.		all at time.
	Pressure at sea level.	Dry bulb.	Wet bulb.	Maximum.	Minimum.	Dew-point.	Relative humidity.	Prevailing direction.	Force.	Average cloudiness.	Maximum.	Minimum.	Total rainfall at m., local time.
1		† 68 70 65 62 70 65 63 63 63 63 63 64 56 58 59 67 70 64 65 64 65 71 64 65 71 64 65 65 71 65 65 71 65 65 71 72 65 65 71 71 71 71 71 71 71 71 71 71 71 71 71	+ 63 63 62 65 64 64.5 65.5 65.5 65.5 67 61.5 63.5 63.5 63.5 63.5 63.5 63.5 63.5 63	74 77 78 73 73 73 73 75 77 77 74 75 77 77 77 77 77 77 77 77 77 77 77 77	64 67 68 63 65 66 65 66 62 62 62 62 64 67 64 64 67 64 64 67 64 64 67 64 65 66 66 67 67 68 68 68 68 68 68 68 68 68 68 68 68 68		60 787 785 785 801 882 576 675 775 687 772 697 697 697 697 697 697 697 697 697 697	ne. ne. ne. s-sw. s-w. nne. nne. nne. nne. w-n. nw-n. ne. ne. ne. ne. sw. s-se-ne. ne. sw-sw-sw. s-se-ne. nne.	\$ 5 4 4 - 2 1 2 4 - 2 1 2 4 - 2 1 2 4 - 2 1 2 2 2 2 2 2 2 2 2 2 2 2 3 3 3 3 3 3	8 2 100 2 110 10 8 4 7 6 6 0 1 - 3 8 7 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 0 1 - 8 8 3 2 - 5	30.12	28. 98 29. 99 29. 97 29. 85 29. 85 29. 85 29. 92 29. 74 29. 78 29. 93 20. 99 20. 99 20. 99 20. 99 20. 99 20. 97 22. 97 22. 97 23. 99 24. 97 25. 97 26. 97 27. 97 28. 97 28	0.58 0.00 0.00 0.00 0.00 0.00 0.00 0.00
ture	+.04	·····			••••	-2.3	-5. 0		·····	-1.2			-4.86

Mean temperature for February, 1900 $(6+2+9)+8=70.5^\circ$; nermal is 70.6°. Mean pressure for January (9+3)+2 is 29.991; normal is 29.949.
*This pressure is as recorded at 1 p. m., Greenwich time. are observed at 6 a. m., local, or 7:29 p. m., Greenwich time. \$\dagger\$ These values are the means of (6+9+2+9)+4. § Beaufort scale.

KITE OBSERVATIONS AT BAYONNE, N. J.

By the Bayonne Kite Club.

The secretary of the Bayonne kite corps, under date of February 19, submits the accompanying table showing the thermometric records and other data accumulated by the corps during the past six months, in continuation of the record published in the Monthly Weather Review for June, 1899, p. 251. The columns 12 to 15 here given were compiled by the Records Division. The altitudes given in the 5th column show a decided gain in the heights from which records are obtained. In ascension No. 118 the record for 1,000 feet is given hourly beside the record for 2,000 feet made by a second thermometer. This was accomplished by means of the kite line transit carrier car. A record was also kept of the electrical phenomena on the kite wire.

The secretary of the club says: